



NATIONAL SENIOR CERTIFICATE EXAMINATION  
NOVEMBER 2022

**PHYSICAL SCIENCES: PAPER II**

**MARKING GUIDELINES**

Time: 3 hours

200 marks

---

**These marking guidelines are prepared for use by examiners and sub-examiners, all of whom are required to attend a standardisation meeting to ensure that the guidelines are consistently interpreted and applied in the marking of candidates' scripts.**

**The IEB will not enter into any discussions or correspondence about any marking guidelines. It is acknowledged that there may be different views about some matters of emphasis or detail in the guidelines. It is also recognised that, without the benefit of attendance at a standardisation meeting, there may be different interpretations of the application of the marking guidelines.**

---

**QUESTION 1         MULTIPLE CHOICE**

- 1.1    D  
 1.2    B  
 1.3    A  
 1.4    C  
 1.5    C  
 1.6    B  
 1.7    D  
 1.8    B  
 1.9    A  
 1.10  D

**QUESTION 2**

- 2.1    2.1.1 An unequal sharing of electrons leading to a dipole forming (as a result of a difference in electronegativity)
- 2.1.2 H-C
- 2.1.3 smallest difference in electronegativity
- 2.2    2.2.1 H must be bonded to:         A small atom  
    With high electronegativity  
    And at least one lone pair of electrons
- 2.2.2 The -O-H bond is very polar/strong dipole forms/H-nucleus is exposed/  
 higher charge density on the hydrogen/ $\delta^+$  and  $\delta^-$  are big (due to large difference in electronegativity)  
 The molecules can get close together/the forces act over shorter distances (due to small atom)
- 2.3    physical intermolecular forces are overcome
- 2.4    Water forms 4 H-bonds with neighbouring molecules.  
 The other two compounds form 2 hydrogen bonds with neighbouring molecules.  
 More energy is needed to overcome the greater number of H-bonds/the stronger IMFs in water (and hence to separate water molecules.)
- 2.5    2.5.1 London/dispersion/induced dipole (forces)
- 2.5.2 **ACCEPT TWO OF:**  
 Ethanol has a larger electron cloud/more electrons  
 Ethanol has a larger interacting (contact) surface/longer chain than methanol  
 Ethanol forms larger/more temporary dipoles

**QUESTION 3**

3.1 3.1.1  $n(\text{Cu}) = \frac{m}{M} = \frac{2,54}{63,5} = \mathbf{0,04 \text{ mol}}$

3.1.2  $n(\text{HNO}_3) = cV = 0,1 \times 0,8 = \mathbf{0,08 \text{ mol}}$

3.1.3  $0,08 \text{ mol HNO}_3 \text{ reacts with } 0,08 \times \frac{3}{8} = 0,03 \text{ mol Cu} < 0,04 \text{ mol}$


**OR:**  $0,04 \text{ mol Cu needs } 0,04 \times \frac{8}{3} = 0,107 \text{ mol HNO}_3 > 0,08 \text{ mol}$

3.1.4  $\mathbf{0,08 \text{ mol HNO}_3}$  (LR) produces  $\frac{0,08}{4} = 0,02 \text{ mol NO}$

Carry over error from 3.1.2 (must use given LR i.e. HNO<sub>3</sub>)

$V(\text{NO}) = nV_m = 0,02 \times 22,4 = \mathbf{0,448 \text{ dm}^3}$

3.2



B: half volume of A  
Gradient less steep

C: same final volume as A  
Steeper gradient than A


- 3.3 More particles per unit volume  
 $\therefore$  more collisions (per unit time)  
 $\therefore$  more effective collisions per unit time  
 $\therefore$  rate will increase

- 3.4 3.4.1 pH will INCREASE.  
 The concentration of acid / HNO<sub>3</sub> / H<sub>3</sub>O<sup>+</sup> / H<sup>+</sup> is **decreasing**  
 as the reaction proceeds.

- 3.4.2 % transmission will DECREASE.  
 (The concentration of Cu(NO<sub>3</sub>)<sub>2</sub> is increasing),  
 so the solution becomes **darker blue**  
 so less light can pass through

- 3.5 3.5.1 A reaction that involves the transfer of electrons

3.5.2 Cu or copper




**QUESTION 4**

4.1 4.1.1 higher than

4.1.2 equal to

4.2 (The mole ratio shows that) for each mole of N<sub>2</sub>O<sub>4</sub> that reacts, two moles of NO<sub>2</sub> are produced

4.3



Rate of reverse (only) increases at t<sub>3</sub>  
 Rate of forward increases and reverse decreases  
 Equilibrium re-established (rates equal and higher than before)  
 –1 if both rates increase together at t<sub>3</sub>


4.4 4.4.1 Both concentrations would sharply decrease at t<sub>4</sub>

4.4.2 since  $c = n / V$  If  $V$  increases,  $c$  must decrease

4.5 in order to decrease the temperature/absorb heat/relieve the stress  
 the forward reaction is favoured  
 because it is endothermic (and absorbs heat)  
 thus the [NO<sub>2</sub>] increases and the [N<sub>2</sub>O<sub>4</sub>] decreases

4.6 The concentrations remain constant/There is no effect.  
 Because:  
 the rate at which N<sub>2</sub>O<sub>4</sub> produces NO<sub>2</sub> and the rate at which NO<sub>2</sub> forms N<sub>2</sub>O<sub>4</sub>  
 increase by the same amount  
 OR the rates of both (the forward and reverse) reactions increase equally/  
 by the same amount

4.7 4.7.1 
$$K_c = \frac{[NO_2]^2}{[N_2O_4]}$$



Initial	0,46	0
Change	- 0,095	+ 0,19
Equilibrium	0,365	0,19

$$K_c = \frac{[0,19]^2}{[0,365]} \text{ coe} = \mathbf{0,099} \text{ OR } 0,10$$

**QUESTION 5**

5.1 5.1.1 proton donor

5.1.2 sulfurous acid (accept sulphurous acid)


5.1.3 an acid that only ionizes partially in an aqueous solution

5.1.4  $\text{H}_3\text{O}^+$  or  $\text{H}^+$  concentration OR  $[\text{H}_3\text{O}^+]$  OR  $[\text{H}^+]$ 

5.1.5 A

5.1.6 EITHER: **B** ( $\text{H}_2\text{SO}_4$ ) is the **strongest** acid solution.  
OR: **C** is the **most concentrated** acid solution.

5.2 5.2.1 amphiprotic OR amphoteric OR ampholyte



pairs are correct labels are correct

5.3 5.3.1  $\text{NH}_4^+ + \text{H}_2\text{O} \rightleftharpoons \text{NH}_3 + \text{H}_3\text{O}^+$ 5.3.2  $\text{F}^- + \text{H}_2\text{O} \rightleftharpoons \text{HF} + \text{OH}^-$ 

(-1 if no reversible arrows – penalise once only)

5.3.3 (a) acidic

(b)  $K_a$  for  $\text{NH}_4^+$ (c)  $\text{NH}_4^+$  ionises more than  $\text{F}^-$   
 $[\text{H}_3\text{O}^+] > [\text{OH}^-]$ 5.4 5.4.1  $\text{Na}_2\text{CO}_3 + 2\text{HBr} \rightarrow 2\text{NaBr} + \text{CO}_2 + \text{H}_2\text{O}$   
products balancing5.4.2  $n(\text{Na}_2\text{CO}_3) = cV = 0,12 \times 0,02 = 0,0024 \text{ mol}$  $n(\text{HBr}) = 2 \times 0,0024 = 0,0048 \text{ mol}$  (2:1) **indicate** ratio used (coe from 5.4.1) $c(\text{HBr}) = n / V = 0,0048 / 0,0152 = 0,3158 \text{ mol.dm}^{-3}$  ✓ (4 d.p.)OR USE:  $\frac{c_a V_a}{a} = \frac{c_b V_b}{b}$  or other version thereof

$$\frac{c_a \times 0,0152}{2} = \frac{0,12 \times 0,0200}{1} \text{ ratio indicated}$$

(coe from 5.4.1)

 $c(\text{HBr}) = 0,3158 \text{ mol.dm}^{-3}$  (4 d.p.)



**QUESTION 7**

7.1 7.1.1 it is inert

7.1.2 positive

7.1.3 reduction

EITHER: electrons are gained

OR: the oxidation number of H decreases

OR: it takes place at the cathode

$$7.1.4 \quad E_{\text{cell}}^{\theta} = E_{\text{cathode}}^{\theta} - E_{\text{anode}}^{\theta} = -0,83 - 1,23 = -2,06 \text{ V}$$

no carry-over

7.1.5 non-spontaneous;  $E_{\text{cell}}^{\theta} < 0$  (must follow from 7.1.4)

7.1.6 (a) blue

(b) blue

7.1.7 (a)  $q = It = 0,05 \times (1,5 \times 3\,600) = 270 \text{ C}$

$$(b) \quad n(e^{-}) = \frac{q}{F} = \frac{270(\text{coe})}{96500} = 0,0028 \text{ mol} \quad \div 96\,500$$

$$[\text{OR } N(e^{-}) = 270 / 1,6 \times 10^{-19} \text{ Then } n(e^{-}) = N / N_A = 0,0028 \text{ mol}]$$

$$n(\text{O}_2) = \frac{0,0028}{4} = 0,0007 \text{ mol} \quad \div 4$$

$$V(\text{O}_2) = nV_m = 0,0007 \times 22,4 \quad \times 22,4$$

$$= 0,016 \text{ dm}^3 \text{ OR } 0,02 \text{ dm}^3$$

7.2 7.2.1 brine

7.2.2 mercury cell.

NaOH produced in separate vessel/area.

7.2.3 diaphragm cell

Contaminant is NaCl

because diaphragm allows anions/ $\text{Cl}^{-}$ /all ions through (non-selective).

**QUESTION 8**

8.1 1-bromo propane

8.2 haloalkane

8.3 8.3.1  $\text{CH}_3\text{CH}=\text{CH}_2$  (–1 if molecular formula given  $\text{C}_3\text{H}_6$ )

8.3.2  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$  propanol -OH on  $\text{C}_1$   
(–1 if molecular formula given  $\text{C}_3\text{H}_8\text{O}$  or  $\text{C}_3\text{H}_7\text{OH}$ )

(–1 in total if structural formula used in 8.3)

8.4 Elimination


8.5 Hydrolysis

8.6 Dehydration

8.7 The organic compounds may have low boiling points and escape from the reaction vessel.



**QUESTION 9**



9.1.2 ester group (COOC) circled (accept COO – as in SAGS)

9.1.3 condensation

9.1.4 ethanoic acid

9.2 heptane-3,3-diol correct chain length diol correct correct numbering  
 mistakes in format or **e** or **ane** omitted (-1)

9.3 9.3.1 (thermal) cracking

9.3.2 prop ene

9.4 9.4.1 Butane: (red-brown) colour remains /(red-brown) colour fades slowly /nothing happens


But-1-ene: (red-brown) colour disappears quickly / immediately

Colour disappears faster for but-1-ene than for butane

1 mark for colour disappearing


2 marks for relative rate of disappearance

9.4.2 substitution



9.4.4 (a)  $\Delta H = E_a - E_{\text{out}} = 4\,795 - 4\,889 = -94 \text{ kJ}\cdot\text{mol}^{-1}$

(b&c)



Max 1/3 if (a) correct  
 but endo curve drawn.  
 Max 2/3 if (a) incorrect  
 and endo curve  
 drawn.

**Total: 200 marks**